



## The Geometric Realization of a Semi-Simplicial Complex

John Milnor

*The Annals of Mathematics*, 2nd Ser., Vol. 65, No. 2. (Mar., 1957), pp. 357-362.

Stable URL:

<http://links.jstor.org/sici?sici=0003-486X%28195703%292%3A65%3A2%3C357%3ATGROAS%3E2.0.CO%3B2-5>

*The Annals of Mathematics* is currently published by Annals of Mathematics.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/annals.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## THE GEOMETRIC REALIZATION OF A SEMI-SIMPLICIAL COMPLEX

BY JOHN MILNOR

(Received February 9, 1956)

Corresponding to each (complete) semi-simplicial complex  $K$ , a topological space  $|K|$  will be defined. This construction will be different from that used by Giever [4] and Hu [5] in that the degeneracy operations of  $K$  are used. This difference is important when dealing with product complexes.

If  $K$  and  $K'$  are countable it is shown that  $|K \times K'|$  is canonically homeomorphic to  $|K| \times |K'|$ . It follows that if  $K$  is a countable group complex then  $|K|$  is a topological group. In particular  $|K(\pi, n)|$  is an abelian topological group.

In the last section it is shown that the space  $|K|$  has the correct singular homology and homotopy groups.

The terminology for semi-simplicial complexes will follow John Moore [7]. In particular the face and degeneracy maps of  $K$  will be denoted by  $\partial_i: K_n \rightarrow K_{n-1}$  and  $s_i: K_n \rightarrow K_{n+1}$  respectively.

### 1. The definition

As standard  $n$ -simplex  $\Delta_n$  take the set of all  $(n + 2)$ -tuples  $(t_0, \dots, t_{n+1})$  satisfying  $0 = t_0 \leq t_1 \leq \dots \leq t_{n+1} = 1$ . The face and degeneracy maps

$$\partial_i: \Delta_{n-1} \rightarrow \Delta_n$$

and  $s_i: \Delta_{n+1} \rightarrow \Delta_n$  are defined by

$$\partial_i(t_0, \dots, t_n) = (t_0, \dots, t_i, t_i, \dots, t_n)$$

$$s_i(t_0, \dots, t_{n+2}) = (t_0, \dots, t_i, t_{i+2}, \dots, t_{n+2}).$$

Let  $K = \bigcup_{i \geq 0} K_i$  be a semi-simplicial complex. Giving  $K$  the discrete topology, form the topological sum

$$\bar{K} = (K_0 \times \Delta_0) + (K_1 \times \Delta_1) + \dots + (K_n \times \Delta_n) + \dots$$

Thus  $\bar{K}$  is a disjoint union of open sets  $k_i \times \Delta_i$ . An equivalence relation in  $\bar{K}$  is generated by the relations

$$(\partial_i k_n, \delta_{n-1}) \sim (k_n, \partial_i \delta_{n-1})$$

$$(s_i k_n, \delta_{n+1}) \sim (k_n, s_i \delta_{n+1}),$$

for each  $k_n \in K_n$ ,  $\delta_{n \pm 1} \in \Delta_{n \pm 1}$  and for  $i = 0, 1, \dots, n$ . The identification space  $|K| = \bar{K}/(\sim)$  will be called the *geometric realization* of  $K$ . The equivalence class of  $(k_n, \delta_n)$  will be denoted by  $|k_n, \delta_n|$ . (The equivalence class  $|k_0, \delta_0|$  may be abbreviated by  $|k_0|$ .)

**THEOREM 1.**  $|K|$  is a CW-complex having one  $n$ -cell corresponding to each non-degenerate  $n$ -simplex of  $K$ .

For the definition of CW-complex see Whitehead [8].

**LEMMA 1.** Every simplex  $k_n \in K_n$  can be expressed in one and only one way as  $k_n = s_{j_p} \cdots s_{j_1} k_{n-p}$  where  $k_{n-p}$  is non-degenerate and  $0 \leq j_1 < \cdots < j_p < n$ . The indices  $j_\alpha$  which occur are precisely those  $j$  for which  $k_n \in s_j K_{n-1}$ .

The proof is not difficult. (See [3] 8.3). Similarly we have:

**LEMMA 2.** Every  $\delta_n \in \Delta_n$  can be written in exactly one way as  $\delta_n = \partial_{i_q} \cdots \partial_{i_1} \delta_{n-q}$  where  $\delta_{n-q}$  is an interior point (that is the coordinates  $t_i$  of  $\delta_{n-q}$  satisfy  $t_0 < t_1 < \cdots < t_{n-q+1}$ ) and  $0 \leq i_1 < \cdots < i_q \leq n$ .

By a non-degenerate point of  $\bar{K}$  will be meant a point  $(k_n, \delta_n)$  with  $k_n$  non-degenerate and  $\delta_n$  interior.

**LEMMA 3.** Each  $(k_n, \delta_n) \in \bar{K}$  is equivalent to a unique non-degenerate point.

Define the map  $\lambda: \bar{K} \rightarrow \bar{K}$  as follows. Given  $k_n$  choose  $j_1, \cdots, j_p, k_{n-p}$  as in Lemma 1 and set

$$\lambda(k_n, \delta_n) = (k_{n-p}, s_{j_1} \cdots s_{j_p} \delta_n).$$

Define the discontinuous function  $\rho: \bar{K} \rightarrow \bar{K}$  by choosing  $i_1 \cdots i_q, \delta_{n-q}$  as in Lemma 2 and setting

$$\rho(k_n, \delta_n) = (\partial_{i_1} \cdots \partial_{i_q} k_n, \delta_{n-q}).$$

Now the composition  $\lambda\rho: \bar{K} \rightarrow \bar{K}$  carries each point into an equivalent, non-degenerate point. It can be verified that if  $x \sim x'$  then  $\lambda\rho(x) = \lambda\rho(x')$ ; which proves Lemma 3.

Take as  $n$ -cells of  $|K|$  the images of the non-degenerate simplexes of  $\bar{K}$ . By Lemma 3 the interiors of these cells partition  $|K|$ . Since the remaining conditions for a CW-complex are easily verified, this proves Theorem 1.

**LEMMA 4.** A semi-simplicial map  $f: K \rightarrow K'$  induces a continuous map  $|K| \rightarrow |K'|$ .

In fact the map  $|f|$  defined by  $|k_n, \delta_n| \rightarrow |f(k_n), \delta_n|$  is clearly well defined and continuous.

As an example of the geometric realization, let  $C$  be an ordered simplicial complex with space  $|C|$ . (See [2] pp. 56 and 67). From  $C$  we can define a semi-simplicial complex  $K$ , where  $K_n$  is the set of all  $(n + 1)$ -tuples  $(a_0, \cdots, a_n)$  of vertices of  $C$  which (1) all lie in a common simplex, and (2) satisfy  $a_0 \leq a_1 \leq \cdots \leq a_n$ . The operations  $\partial_i, s_i$  are defined in the usual way.

**ASSERTION.** The space  $|C|$  is homeomorphic to the geometric realization  $|K|$ . In fact the point  $|(a_0, \cdots, a_n); (t_0, \cdots, t_{n+1})|$  of  $|K|$  corresponds to the point of  $|C|$  whose  $a^{\text{th}}$  barycentric coordinate,  $a$  being a vertex of  $C$ , is the sum, over all  $i$  for which  $a_i = a$ , of  $t_{i+1} - t_i$ . The proof is easily given.

## 2. Product complexes

Let  $K \times K'$  be the cartesian product of two semi-simplicial complexes (that is  $(K \times K')_n = K_n \times K'_n$ ). The projection maps  $\rho: K \times K' \rightarrow K$  and  $\rho': K \times K' \rightarrow K'$  induce maps  $|\rho|$  and  $|\rho'|$  of the geometric realizations. A map

$$\eta: |K \times K'| \rightarrow |K| \times |K'|$$

is defined by  $\eta = |\rho| \times |\rho'|$ .

**THEOREM 2.**  $\eta$  is a one-one map of  $|K \times K'|$  onto  $|K| \times |K'|$ . If either (a)  $K$  and  $K'$  are countable, or (b) one of the two CW-complexes  $|K|$ ,  $|K'|$  is locally finite; then  $\eta$  is a homeomorphism.

The restrictions (a) or (b) are necessary in order to prove that  $|K| \times |K'|$  is a CW-complex. (For the proof in case (b) see [8] p. 227 and for case (a) see [6] 2.1.)

**PROOF** (Compare [2] p. 68). If  $x''$  is a point of  $|K \times K'|$  with non-degenerate representative  $(k_n \times k'_n, \delta_n)$  we will first determine the non-degenerate representative of  $|\rho|(x'') = |k_n, \delta_n|$ . Since  $\delta_n$  is an interior point of  $\Delta_n$ , this representative has the form

$$(k_{n-p}, s_{i_1} \cdots s_{i_p} \delta_n) \quad \text{where} \quad k_n = s_{i_p} \cdots s_{i_1} k_{n-p}$$

(see proof of Lemma 3). Similarly  $|\rho'|(x'')$  is represented by

$$(k'_{n-q}, s_{j_1} \cdots s_{j_q} \delta_n)$$

where  $k'_n = s_{j_q} \cdots s_{j_1} k'_{n-q}$ . The indices  $i_\alpha$  and  $j_\beta$  must be distinct; for if  $i_\alpha = j_\beta$  for some  $\alpha, \beta$  then  $k_n \times k'_n$  would be an element of  $s_{i_\alpha}(K_{n-1} \times K'_{n-1})$ .

However the point  $x''$  can be completely determined by its image.

$$|k_{n-p}, s_{i_1} \cdots s_{i_p} \delta_n| \times |k'_{n-q}, s_{j_1} \cdots s_{j_q} \delta_n|.$$

In fact given any pair  $(x, x') \in |K| \times |K'|$  define  $\bar{\eta}(x, x') \in |K \times K'|$  as follows. Let  $(k_a, \delta_a)$  and  $(k'_b, \delta'_b)$  be the non-degenerate representatives: where  $\delta_a = (t_0, \dots, t_{a+1})$ ,  $\delta'_b = (u_0, \dots, u_{b+1})$ . Let  $0 = w_0 < \dots < w_{n+1} = 1$  be the distinct numbers  $t_i$  and  $u_j$  arranged in order. Set  $\delta''_n = (w_0, \dots, w_{n+1})$ . Then if  $\mu_1 < \dots < \mu_{n-a}$  are those integers  $\mu = 0, 1, \dots, n-1$  such that  $w_{\mu+1}$  is not one of the  $t_i$ , we have  $\delta_a = s_{\mu_1} \cdots s_{\mu_{n-a}} \delta''_n$ . Similarly  $\delta'_b = s_{\nu_1} \cdots s_{\nu_{n-b}} \delta''_n$  where the sets  $\{\mu_i\}$  and  $\{\nu_j\}$  are disjoint. Now define

$$\bar{\eta}(x, x') = |(s_{\mu_{n-a}} \cdots s_{\mu_1} k_a) \times (s_{\nu_{n-b}} \cdots s_{\nu_1} k'_b), \delta''_n|.$$

Clearly

$$\begin{aligned} |\rho| \bar{\eta}(x, x') &= |s_{\mu_{n-a}} \cdots s_{\mu_1} k_a, \delta''_n| = |k_a, s_{\mu_1} \cdots s_{\mu_{n-a}} \delta''_n| \\ &= |k_a, \delta_a| = x \end{aligned}$$

and  $|\rho'| \bar{\eta}(x, x') = x'$ , which proves that  $\eta \bar{\eta}$  is the identity map of  $|K| \times |K'|$ . On the other hand, taking  $x''$  as above we have

$$\begin{aligned} \bar{\eta}(x'') &= \bar{\eta}(|k_{n-p}, s_{i_1} \cdots s_{i_p} \delta_n|, |k'_{n-q}, s_{j_1} \cdots s_{j_q} \delta_n|) \\ &= |(s_{i_p} \cdots s_{i_1} k_{n-p}) \times (s_{j_q} \cdots s_{j_1} k'_{n-q}), \delta_n| = x''. \end{aligned}$$

To complete the proof it is only necessary to show that  $\bar{\eta}$  is continuous. However it is easily verified that  $\bar{\eta}$  is continuous on each product cell of  $|K| \times |K'|$ . Since we know that this product is a CW-complex, this completes the proof.

An important special case is the following. Let  $I$  denote the semi-simplicial complex consisting of a 1-simplex and its faces and degeneracies.

**COROLLARY.** *A semi-simplicial homotopy  $h:K \times I \rightarrow K'$  induces an ordinary homotopy  $|K| \times [0, 1] \rightarrow |K'|$ .*

In fact the interval  $[0, 1]$  may be identified with  $|I|$ . The homotopy is now given by the composition

$$|K| \times |I| \xrightarrow{\bar{\eta}} |K \times I| \xrightarrow{|h|} |K'|.$$

### 3. Product operations

Now let  $K$  be a countable complex. Any semi-simplicial map  $p:K \times K \rightarrow K$  induces by Lemma 4 and Theorem 2 a continuous product

$$|p| \bar{\eta}: |K| \times |K| \rightarrow |K|.$$

If there is an element  $e_0$  in  $K_0$  such that  $s_0^n e_0$  is a two-sided identity in  $K_n$  for each  $n$ , then it follows that  $|e_0|$  is a two-sided identity in  $|K|$ ; so that  $|K|$  is an  $H$ -space. If the product operation  $p$  is associative or commutative then it is easily verified that  $|p| \bar{\eta}$  is associative or commutative. Hence we have the following.

**THEOREM 3.** *If  $K$  is a countable group complex (countable abelian group complex), then  $|K|$  is a topological group (abelian topological group).*

Let  $K(\pi, n)$  denote the Eilenberg MacLane semi-simplicial complex (see [1]). Since  $K(\pi, n)$  is an abelian group complex we have:

**COROLLARY.** *If  $\pi$  is a countable abelian group, then for  $n \geq 0$  the geometric realization  $|K(\pi, n)|$  is an abelian topological group.*

It will be shown in the next section that  $|K(\pi, n)|$  actually is a space with one non-vanishing homotopy group.

The above construction can also be applied to other algebraic operations. For example a pairing  $K \times K' \rightarrow K''$  between countable group complexes induces a pairing between their realizations. If  $K$  is a countable semi-simplicial complex of  $\Lambda$ -modules, where  $\Lambda$  is a discrete ring, then  $|K|$  is a topological  $\Lambda$ -module.

### 4. The topology of $|K|$

For any space  $X$  let  $S(X)$  be the total singular complex. For any semi-simplicial complex  $K$  a one-one semi-simplicial map  $i:K \rightarrow S(|K|)$  is defined by

$$i(k_n)(\delta_n) = |k_n, \delta_n|.$$

Let  $H_*(K)$  denote homology with integer coefficients.

**LEMMA 5.** *The inclusion  $K \rightarrow S(|K|)$  induces an isomorphism  $H_*(K) \cong H_*(S|K|)$  of homology groups.*

By the  $n$ -skeleton  $K^{(n)}$  of  $K$  is meant the subcomplex consisting of all  $K_i$ ,  $i \leq n$  and their degeneracies. Thus  $|K^{(n)}|$  is just the  $n$ -skeleton of  $|K|$  considered as a  $CW$ -complex. The sequence of subcomplexes

$$K^{(0)} \subset K^{(1)} \subset \dots$$

gives rise to a spectral sequence  $\{E_{pq}^r\}$ ; where  $E^\infty$  is the graded group corresponding to  $H_*(K)$  under the induced filtration; and

$$E_{pq}^1 = H_{p+q}(K^{(p)} \text{ mod } K^{(p-1)}).$$

It is easily verified that  $E_{pq}^1 = 0$  for  $q \neq 0$ , and that  $E_{p0}^1$  is the free abelian group generated by the non-degenerate  $p$ -simplexes of  $K$ . From the first assertion it follows that  $E_{p0}^2 = E_{p0}^\infty = H_p(K)$ .

On the other hand the sequence

$$S(|K^{(0)}|) \subset S(|K^{(1)}|) \subset \dots$$

gives rise to a spectral sequence  $\{\bar{E}_{pq}^r\}$  where  $\bar{E}^\infty$  is the graded group corresponding to  $H_*(S(|K|))$ . Since it is easily verified that the induced map  $E_{pq}^1 \rightarrow \bar{E}_{pq}^1$  is an isomorphism, it follows that the rest of the spectral sequence is also mapped isomorphically; which completes the proof.

Now suppose that  $K$  satisfies the Kan extension condition, so that  $\pi_1(K, k_0)$  can be defined.

LEMMA 6. *If  $K$  is a Kan complex then the inclusion  $i$  induces an isomorphism of  $\pi_1(K, k_0)$  onto  $\pi_1(S(|K|), i(k_0)) = \pi_1(|K|, |k_0|)$ .*

Let  $K'$  be the Eilenberg subcomplex consisting of those simplices of  $K$  whose vertices are all at  $k_0$ . Then  $\pi_1(K, k_0)$  can be considered as a group with one generator for each element of  $K'_1$  and one relation for each element of  $K'_2$ .

The space  $|K'|$  is a  $CW$ -complex with one vertex. For such a space the group  $\pi_1$  is known to have one generator for each edge and one relation for each face. Comparing these two descriptions it follows easily that the homomorphism  $\pi_1(K) = \pi_1(K') \rightarrow \pi_1(|K'|)$  is an isomorphism.

We may assume that  $K$  is connected. Then it is known (see [7] Chapter I, appendix C) that the inclusion map  $K' \rightarrow K$  is a semi-simplicial homotopy equivalence. By the corollary to Theorem 2 this proves that the inclusion  $|K'| \rightarrow |K|$  is a homotopy equivalence; which completes the proof of Lemma 6.

REMARK 1. From Lemmas 5 and 6 it can be proved, using a relative Hurewicz theorem, that the homomorphisms

$$\pi_n(K, k_0) \rightarrow \pi_n(|K|, |k_0|)$$

are isomorphisms for all  $n$ . (The proof of the relative Hurewicz theorem given in [9] §3 carries over to the semi-simplicial case without essential change, making use of [7] Chapter I, appendices A and C. This theorem is applied to the pair  $(S(|\bar{K}|), \bar{K})$  where  $\bar{K}$  denotes the universal covering complex of  $K$ .)

REMARK 2. The space  $|K(\pi, n)|$  has  $n^{\text{th}}$  homotopy group  $\pi$ , and other homotopy groups trivial. This clearly follows from the preceding remark. Alternatively the proof given by Hu [5] may be used without essential change.

Now let  $X$  be any topological space. There is a canonical map

$$j: |S(X)| \rightarrow X$$

defined by  $j(|k_n, \delta_n|) = k_n(\delta_n)$ .

**THEOREM 4.** *The map  $j: |S(X)| \rightarrow X$  induces isomorphisms of the singular homology and homotopy groups.*

(This result is essentially due to Giever [4]).

The map  $j$  induces a semi-simplicial map  $j_*: S(|S(X)|) \rightarrow S(X)$ . A map  $i$  in the opposite direction was defined at the beginning of this section. The composition  $j_*i: S(X) \rightarrow S(X)$  is the identity map. Together with Lemma 5 this implies that  $j$  induces isomorphisms of the singular homology groups of  $|S(X)|$  onto those of  $X$ . Together with Remark 1 it implies that  $j$  induces isomorphisms of the homotopy groups of  $|S(X)|$  onto those of  $X$ . This completes the proof.

PRINCETON UNIVERSITY

#### REFERENCES

1. S. EILENBERG and S. MACLANE, *Relations between homology and homotopy groups of spaces II*, Ann. of Math, 51 (1950), 514-533.
2. ——— and N. STEENROD, *Foundations of Algebraic Topology*, Princeton, 1952.
3. ——— and J. A. ZILBER, *Semi-simplicial complexes and singular homology*, Ann. of Math., 51 (1950), 499-513.
4. J. B. GIEVER, *On the equivalence of two singular homology theories*, Ann. of Math., 51 (1950), 178-191.
5. S. T. HU, *On the realizability of homotopy groups and their operations*, Pacific J. Math., 1 (1951), 583-602.
6. J. MILNOR, *Construction of universal bundles I*, Ann. of Math., 63 (1956), 272-284.
7. J. MOORE, *Algebraic homotopy theory (Lecture notes)*, Princeton, 1955-56.
8. J. H. C. WHITEHEAD, *Combinatorial homotopy I*, Bull. Amer. Math. Soc., 55 (1949), 213-245.
9. J. MOORE, *Some applications of homology theory to homotopy problems*, Ann. of Math., 58 (1953), 325-350.